

VU Research Portal

Genome-wide association study (GWAS)-identified disease risk alleles do not compromise human longevity

Beekman, M.; Nederstigt, C.; Suchiman, H.E.D.; Kremer, D.; van der Breggen, R.; Lakenberg, N.; Alemayehu, W.G.; de Craen, A.J.; Westendorp, R.G.J.; Boomsma, D.I.; de Geus, E.J.C.; Houwing-Duistermaat, J.J.; Heijmans, B.T.; Slagboom, P.E.

published in

Proceedings of the National Academy of Sciences of the United States of America
2010

DOI (link to publisher)

[10.1073/pnas.1003540107](https://doi.org/10.1073/pnas.1003540107)

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Beekman, M., Nederstigt, C., Suchiman, H. E. D., Kremer, D., van der Breggen, R., Lakenberg, N., Alemayehu, W. G., de Craen, A. J., Westendorp, R. G. J., Boomsma, D. I., de Geus, E. J. C., Houwing-Duistermaat, J. J., Heijmans, B. T., & Slagboom, P. E. (2010). Genome-wide association study (GWAS)-identified disease risk alleles do not compromise human longevity. *Proceedings of the National Academy of Sciences of the United States of America*, 107(42), 18046-18049. <https://doi.org/10.1073/pnas.1003540107>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Supporting Information

Beekman et al. 10.1073/pnas.1003540107




Fig. S1. The study population was divided into 10 age groups: (1) <26 y ($n = 116$), (2) 26–35 y ($n = 576$), (3) 36–45 y ($n = 247$), (4) 46–55 y ($n = 344$), (5) 56–65 y ($n = 457$), (6) 66–75 y ($n = 135$), (7) 76–85 y ($n = 547$), (8) 86–95 y ($n = 1,029$), (9) 96–98 y ($n = 101$), and (10) ≥ 99 y ($n = 38$). Per age group, the mean number of risk alleles was determined for the sum of all risk alleles (blue), sum of the subset of metabolic risk alleles (pink), and subset of the cancer risk alleles (yellow).




Fig. S2. Four small clusters are located apart from the main large cluster. These four clusters contain four separate families, each consisting of three nonagenarians. Because these four families report that they are of Dutch ancestry and because these are single-family clusters relatively close to the majority of the samples, we conclude that there is no substructure among the participants of the LLS to an extent that would affect our conclusions.

Table S1. Overview of loci associated with heart disease, cancer, and type 2 diabetes up to February 2009 in at least two independent GWASs

Locus	Chromosome	Nearest genes	SNP*	Position	Risk/Non risk [†]	Associated disease	OR [§]	References
1	1p13.3	CELSR2, PSRC1	rs599839 rs646776	109623689 109620053	A/G T/C	CAD CAD	1.39 —	(1, 2) Surrogate for rs599839 ($r^2 = 0.94$)
2	2q32.3	TMEFF2	rs10497721 rs10497726 rs10497723	192622607 192759565 192817829	A/C[‡] C/A [‡] G/A	T2D CAD T2D	— — 2.32	(3) (4) (5)
3	3p25.2	PPARG	rs17036101 rs1801282	12252845 12368125	G/A C/G	T2D T2D	1.15 1.14	(6) (7–11)
4	3q27.2	IGF2BP2	rs4402960 rs1470579	186994381 187011774	T/G C/A	T2D T2D	1.14 1.14	(7–9, 12) (7, 13)
5	4p16.1	WFS1	rs10010131 rs6446482 rs734312	6343816 6346594 6354255	G/A G/C A/G	T2D T2D T2D	1.15 1.15 1.23/1.25 [¶]	(14, 15) (14, 15) (14, 15)
6	5p15.33	TERT-CLPTM1L	rs402710 rs401681	1373722 1375087	C/T G/A	LC LC	1.18 1.15	(16) (17)
7	5q34	BC011998	rs10515869 rs6556756 rs9314033	163444804 163821858 163822784	G/A[‡] G/T[‡] C/A [‡]	HF BC BC	— — —	(4) (18) (18)
8	6p22.3	CDKAL1	rs10946398 rs7754840 rs7756992 rs9465871	20769013 20769229 20787688 20825234	C/A C/G G/A C/T	T2D T2D T2D T2D	1.12 1.12 1.20 1.18/2.17 [¶]	(9, 19) (7, 8, 20) (12, 21) (10)
9	6q25.1	MTHFD1L	rs6922269	151294678	A/G	CAD	1.37	(1, 10)
10	8q24.11	SLC30A8	rs13266634	118253964	C/T	T2D	1.12	(7–9, 12, 19, 21, 22)
11	8q24.21	POU5F1, LOC727677	rs7001069 rs10505483 rs13281615 rs10505477 rs11985829 rs10808556 rs6983267 rs7013278 rs10505474 rs2060776 rs10956369 rs7014346 rs4871789 rs7842552 rs1447295 rs4242382 rs7837688 rs564398 rs10757274 rs1537371 rs1556516 rs10511701 rs2383206 rs2383207 rs10757278 rs1333049 rs10811661	128179828 128194377 128424800 128476625 128478414 128482329 128482487 128484074 128486686 128489299 128492999 128493974 128497243 128500876 128554220 128586755 128608542 22019547 22086055 22089568 22090176 22102599 22105026 22105959 22114477 22115503 22124094	A/G [‡] G/A [‡] G/A G/A C/T [‡] A/G G/T C/T [‡] G/A [‡] T/G [‡] A/T [‡] A/G A/G [‡] G/A A/C A/G T/G T/C G/A A/C [‡] C/G [‡] C/T [‡] G/A G/A G/A C/G T/C	PC PC BC CC, PC CC, PC CC, PC CC, PC CC CC CC CC CC CC CC CC PC PC PC CAD CVD CVD CVD CAD CAD CAD CAD T2D	— — 1.08 1.27/1.43 1.08/1.22 1.26/1.31 1.25/1.20 — — — — 1.19 — 1.15 1.43/2.23 [¶] 1.66/2.22 [¶] 1.46/2.03 1.12/1.21 [¶] 1.18/1.29 [¶] — — — — 1.26/1.26 [¶] 1.25 1.28 1.47/1.90 [¶] 1.20	(18) (18) (23) (24–26) (27) (26, 27) (25, 27–30) (27) (27) (27) (27) (27, 31) (27) (31) (26, 28, 32, 33) (29) (28) (9, 34) (35) (4) (4) (4) (35) (36) (11, 36) (1, 10) (7–9, 11, 12, 19)
12	9p21.3	CDKN2BAS	rs1111875 rs5015480 rs7923837	94452862 94455539 94471897	C/T C/T A/G	T2D T2D T2D	1.13 1.13 1.22/1.45 [¶]	(7–9, 12, 22) (9) (12, 22)
13	10q23.33	HHEX	rs7901695 rs4506565 rs7903146 rs12255372	114744078 114746031 114748339 114798892	C/T T/A T/C T/G	T2D T2D T2D, CC T2D, BC, PC	1.37 1.36/1.88 [¶] 1.37/1.25–2.15 [¶] 1.64/1.21–1.37 [¶] /1.09–1.15 [¶]	(9, 37) (10) (7, 10, 19, 22, 37–40) (37–39, 41, 42)
14	10q25.2	TCF7L2	rs1219648 rs2420946 rs2981582	123336180 123341314 123342307	G/A T/C A/G	BC BC BC	1.32 1.32 1.26	(43, 44) (43, 45) (23, 45)

Table S1. Cont.

Locus	Chromosome	Nearest genes	SNP*	Position	Risk/Non risk [†]	Associated disease	OR [§]	References
16	11p15.1	KCNJ11	rs5215	17365206	C/T	T2D	1.14	(9)
			rs5219	17366148	T/C	T2D	1.14	(7, 8, 12, 19)
17	12q21.1	TSPAN8	rs1495377	69863368	G/C	T2D	1.28/1.51 [¶]	(10)
			rs7961581	69949369	C/T	T2D	1.09	(6)
18	15q25.1	LOC123688, CHRNA3	rs8034191	76593078	T/C	LC	1.30	(46–48)
			rs1051730	76681394	G/A	LC	1.31	(16, 46, 47, 49)
			rs8042374	76695087	G/A [‡]	LC	—	(17)
19	16q12.1	TOX3	rs8051542	51091668	T/C	BC	1.09	(23)
			rs12443621	51105538	G/A	BC	1.11	(23)
			rs3803662	51143842	T/C	BC	1.20	(23, 50, 51)
20	16q12.2	FTO	rs8050136	52373776	A/C	T2D	1.17	(8, 9, 19, 52)
			rs9939609	52378028	A/T	T2D	1.34/1.55 [¶]	(10, 52)
21	17q12	HNF1B	rs757210	33170628	T/C	T2D	1.12	(53, 54)
			rs4430796	33172153	A/G	PC, T2D	1.22/0.91 [¶]	(29, 54)
			rs7501939	33175269	C/T	PC	1.19	(54, 55)
			rs3760511	33180426	C/A	PC	1.16	(54)
22	18q21.1	SMAD7	rs4939827	44707461	T/C	CC	1.18	(30, 31, 56)
			rs12953717	44707927	T/C	CC	1.17	(31, 56)
			rs4464148	44713030	C/T	CC	1.15	(56)
23	Xp11.22	NUDT11	rs5945572	51246423	A/G	PC	1.23	(57)
			rs5945619	51258412	T/C	PC	1.19	(58)

BC, breast cancer; CAD, coronary artery disease; CC, colon carcinoma; CVD, cardiovascular disease; HF, heart failure; LC, lung cancer; PC, prostate cancer; T2D, type 2 diabetes.

*SNPs genotyped in the LLS are denoted in bold.

[†]Major allele of genotyped SNPs is underlined.

*Risk allele is not found in the literature; data are retrieved from SnpPer (ChIP Bioinformatics), and the minor allele is denoted first.

[§]OR per allele.

¹OR in heterozygotes and homozygotes, respectively.

1. Samani NJ, et al.; WTCCC and the Cardiogenics Consortium (2007) Genomewide association analysis of coronary artery disease. *N Engl J Med* 357:443–453.
2. Willer CJ, et al. (2008) Newly identified loci that influence lipid concentrations and risk of coronary artery disease. *Nat Genet* 40:161–169.
3. Meigs JB, et al. (2007) Genome-wide association with diabetes-related traits in the Framingham Heart Study. *BMC Med Genet* 8(Suppl 1):S16.
4. Larson MG, et al. (2007) Framingham Heart Study 100K project: Genome-wide associations for cardiovascular disease outcomes. *BMC Med Genet* 8(Suppl 1):S5.
5. Hayes MG, et al. (2007) Identification of type 2 diabetes genes in Mexican Americans through genome-wide association studies. *Diabetes* 56:3033–3044.
6. Zeggini E, et al.; Wellcome Trust Case Control Consortium (2008) Meta-analysis of genome-wide association data and large-scale replication identifies additional susceptibility loci for type 2 diabetes. *Nat Genet* 40:638–645.
7. Saxena R, et al.; Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research (2007) Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. *Science* 316:1331–1336.
8. Scott LJ, et al. (2007) A genome-wide association study of type 2 diabetes in Finns detects multiple susceptibility variants. *Science* 316:1341–1345.
9. Zeggini E, et al.; Wellcome Trust Case Control Consortium (WTCCC) (2007) Replication of genome-wide association signals in UK samples reveals risk loci for type 2 diabetes. *Science* 316:1336–1341.
10. Wellcome Trust Case Control Consortium (2007) Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 447:661–678.
11. Helgadottir A, et al. (2008) The same sequence variant on 9p21 associates with myocardial infarction, abdominal aortic aneurysm and intracranial aneurysm. *Nat Genet* 40:217–224.
12. Omori S, et al. (2008) Association of CDKAL1, IGF2BP2, CDKN2A/B, HHEX, SLC30A8, and KCNJ11 with susceptibility to type 2 diabetes in a Japanese population. *Diabetes* 57:791–795.
13. Cauchi S, et al. (2008) Post genome-wide association studies of novel genes associated with type 2 diabetes show gene-gene interaction and high predictive value. *PLoS ONE* 3:e2031.
14. Sandhu MS, et al. (2007) Common variants in WFS1 confer risk of type 2 diabetes. *Nat Genet* 39:951–953.
15. Franks PW, et al. (2008) Replication of the association between variants in WFS1 and risk of type 2 diabetes in European populations. *Diabetologia* 51:458–463.
16. McKay JD, et al.; EPIC Study (2008) Lung cancer susceptibility locus at 5p15.33. *Nat Genet* 40:1404–1406.
17. Wang Y, et al. (2008) Common 5p15.33 and 6p21.33 variants influence lung cancer risk. *Nat Genet* 40:1407–1409.
18. Murabito JM, et al. (2007) A genome-wide association study of breast and prostate cancer in the NHLBI's Framingham Heart Study. *BMC Med Genet* 8(Suppl 1):S6.
19. Timpson NJ, et al. (2009) Adiposity-related heterogeneity in patterns of type 2 diabetes susceptibility observed in genome-wide association data. *Diabetes* 58:505–510.
20. Stancáková A, et al.; EUGENE2 Consortium (2008) Single-nucleotide polymorphism rs7754840 of CDKAL1 is associated with impaired insulin secretion in nondiabetic offspring of type 2 diabetic subjects and in a large sample of men with normal glucose tolerance. *J Clin Endocrinol Metab* 93:1924–1930.
21. Steinthorsdóttir V, et al. (2007) A variant in CDKAL1 influences insulin response and risk of type 2 diabetes. *Nat Genet* 39:770–775.
22. Sladek R, et al. (2007) A genome-wide association study identifies novel risk loci for type 2 diabetes. *Nature* 445:881–885.
23. Easton DF, et al.; SEARCH collaborators; kConFab; AOCs Management Group (2007) Genome-wide association study identifies novel breast cancer susceptibility loci. *Nature* 447:1087–1093.
24. Zanke BW, et al. (2007) Genome-wide association scan identifies a colorectal cancer susceptibility locus on chromosome 8q24. *Nat Genet* 39:989–994.
25. Tomlinson I, et al.; CORGI Consortium (2007) A genome-wide association scan of tag SNPs identifies a susceptibility variant for colorectal cancer at 8q24.21. *Nat Genet* 39:984–988.
26. Ghoussaini M, et al.; UK Genetic Prostate Cancer Study Collaborators/British Association of Urological Surgeons' Section of Oncology; UK ProtecT Study Collaborators (2008) Multiple loci with different cancer specificities within the 8q24 gene desert. *J Natl Cancer Inst* 100:962–966.
27. Haiman CA, et al. (2007) A common genetic risk factor for colorectal and prostate cancer. *Nat Genet* 39:954–956.
28. Yeager M, et al. (2007) Genome-wide association study of prostate cancer identifies a second risk locus at 8q24. *Nat Genet* 39:645–649.
29. Thomas G, et al. (2008) Multiple loci identified in a genome-wide association study of prostate cancer. *Nat Genet* 40:310–315.
30. Tomlinson IP, et al.; CORGI Consortium; EPICOLON Consortium (2008) A genome-wide association study identifies colorectal cancer susceptibility loci on chromosomes 10p14 and 8q23.3. *Nat Genet* 40:623–630.
31. Tenesa A, et al. (2008) Genome-wide association scan identifies a colorectal cancer susceptibility locus on 11q23 and replicates risk loci at 8q24 and 18q21. *Nat Genet* 40:631–637.
32. Severi G, et al. (2007) The common variant rs1447295 on chromosome 8q24 and prostate cancer risk: Results from an Australian population-based case-control study. *Cancer Epidemiol Biomarkers Prev* 16:610–612.
33. Cheng I, et al. (2008) 8q24 and prostate cancer: Association with advanced disease and meta-analysis. *Eur J Hum Genet* 16:496–505.

34. Broadbent HM, et al.; PROCARDIS consortium (2008) Susceptibility to coronary artery disease and diabetes is encoded by distinct, tightly linked SNPs in the ANRIL locus on chromosome 9p. *Hum Mol Genet* 17:806–814.
35. McPherson R, et al. (2007) A common allele on chromosome 9 associated with coronary heart disease. *Science* 316:1488–1491.
36. Helgadottir A, et al. (2007) A common variant on chromosome 9p21 affects the risk of myocardial infarction. *Science* 316:1491–1493.
37. Salonen JT, et al. (2007) Type 2 diabetes whole-genome association study in four populations: The DiaGen consortium. *Am J Hum Genet* 81:338–345.
38. Lysenko V, et al. (2007) Mechanisms by which common variants in the TCF7L2 gene increase risk of type 2 diabetes. *J Clin Invest* 117:2155–2163.
39. Palmer ND, et al. (2008) Association of TCF7L2 gene polymorphisms with reduced acute insulin response in Hispanic Americans. *J Clin Endocrinol Metab* 93:304–309.
40. Folsom AR, et al. (2008) Variation in TCF7L2 and increased risk of colon cancer: The Atherosclerosis Risk in Communities (ARIC) Study. *Diabetes Care* 31:905–909.
41. Burwinkel B, et al. (2006) Transcription factor 7-like 2 (TCF7L2) variant is associated with familial breast cancer risk: A case-control study. *BMC Cancer* 6:268.
42. Agalliu I, et al. (2008) Evaluation of a variant in the transcription factor 7-like 2 (TCF7L2) gene and prostate cancer risk in a population-based study. *Prostate* 68:740–747.
43. Hunter DJ, et al. (2007) A genome-wide association study identifies alleles in FGFR2 associated with risk of sporadic postmenopausal breast cancer. *Nat Genet* 39:870–874.
44. Stacey SN, et al. (2008) Common variants on chromosome 5p12 confer susceptibility to estrogen receptor-positive breast cancer. *Nat Genet* 40:703–706.
45. Raskin L, et al. (2008) FGFR2 is a breast cancer susceptibility gene in Jewish and Arab Israeli populations. *Cancer Epidemiol Biomarkers Prev* 17:1060–1065.
46. Amos CI, et al. (2008) Genome-wide association scan of tag SNPs identifies a susceptibility locus for lung cancer at 15q25.1. *Nat Genet* 40:616–622.
47. Hung RJ, et al. (2008) A susceptibility locus for lung cancer maps to nicotinic acetylcholine receptor subunit genes on 15p25. *Nature* 452:633–637.
48. Liu P, et al. (2008) Familial aggregation of common sequence variants on 15q24-25.1 in lung cancer. *J Natl Cancer Inst* 100:1326–1330.
49. Thorgeirsson TE, et al. (2008) A variant associated with nicotine dependence, lung cancer and peripheral arterial disease. *Nature* 452:638–642.
50. Stacey SN, et al. (2007) Common variants on chromosomes 2q35 and 16q12 confer susceptibility to estrogen receptor-positive breast cancer. *Nat Genet* 39:865–869.
51. Garcia-Closas M, et al.; Australian Ovarian Cancer Management Group; Kathleen Cunningham Foundation Consortium for Research into Familial Breast Cancer (2008) Heterogeneity of breast cancer associations with five susceptibility loci by clinical and pathological characteristics. *PLoS Genet* 4:e1000054.
52. Scuteri A, et al. (2007) Genome-wide association scan shows genetic variants in the FTO gene are associated with obesity-related traits. *PLoS Genet* 3:e115.
53. Winkler W, et al. (2007) Evaluation of common variants in the six known maturity-onset diabetes of the young (MODY) genes for association with type 2 diabetes. *Diabetes* 56:685–693.
54. Gudmundsson J, et al. (2007) Two variants on chromosome 17 confer prostate cancer risk, and the one in TCF2 protects against type 2 diabetes. *Nat Genet* 39:977–983.
55. Levin AM, et al. (2008) Chromosome 17q12 variants contribute to risk of early-onset prostate cancer. *Cancer Res* 68:6492–6495.
56. Broderick P, et al.; CORGI Consortium (2007) A genome-wide association study shows that common alleles of SMAD7 influence colorectal cancer risk. *Nat Genet* 39:1315–1317.
57. Gudmundsson J, et al. (2008) Common sequence variants on 2p15 and Xp11.22 confer susceptibility to prostate cancer. *Nat Genet* 40:281–283.
58. Eeles RA, et al.; UK Genetic Prostate Cancer Study Collaborators; British Association of Urological Surgeons' Section of Oncology; UK ProtecT Study Collaborators (2008) Multiple newly identified loci associated with prostate cancer susceptibility. *Nat Genet* 40:316–321.

Table S2. Cross-tabulation of 10% lower and upper tails of the risk allele distribution in the control and long-lived groups

	All risk alleles	
	<23 alleles (9.4%)	≥31 alleles (12.9%)
Control	172 (9.1%)	237 (12.6%)
Long-lived	166 (9.8%)	225 (13.2%)

OR = 0.99; 95% CI: 0.97–1.03; *P* = 0.909.

Table S3. Cross-tabulation of 10% lower and upper tails of the metabolic risk allele distribution in the control and long-lived groups

	Metabolic risk alleles	
	<15 alleles (6.8%)	≥23 alleles (8.5%)
Control	136 (7.2%)	176 (9.3%)
Long-lived	108 (6.3%)	129 (7.6%)

OR = 0.99; 95% CI: 0.95–1.03; $P = 0.644$.

Table S4. Cross-tabulation of 10% lower and upper tails of the cancer risk allele distribution in the control and long-lived groups

	Cancer risk alleles	
	≤7 alleles (11.9%)	≥13 alleles (12%)
Control	220 (11.7%)	244 (12.9%)
Long-lived	208 (12.2%)	188 (11.0%)

OR = 0.97; 95% CI: 0.92–1.01; $P = 0.135$.

Table S5. Association of GWA-identified disease risk alleles with longevity

					Risk allele frequency				Meta-analysis		
Chromosome	SNP	Position	Disease profile*	Risk/ Nonrisk [‡]	LLS controls	LLS 90+ subjects	NTR controls	Leiden 85 Plus Study subjects	OR [§]	95% CI [§]	P value [¶]
1p13.3	rs646776	109620053	M	T/C	0.786	0.737	0.778	0.777	0.88	0.79–0.99	0.035
2q32.3	rs10497721	192622607	M	A/C	0.092	0.113	0.093	0.090	1.07	0.91–1.26	0.367
3p25.2	rs1801282	12368125	M	C/G	0.884	0.883	0.892	0.873	0.89	0.76–1.03	0.131
3q27.2	rs4402960	186994381	M	T/G	0.316	0.284	0.300	0.305	0.96	0.86–1.06	0.374
4p16.1	rs10010131	6343816	M	G/A	0.586	0.562	0.590	0.600	0.98	0.89–1.08	0.742
5q34	rs10515869	163444804	—	A/G	0.444	0.424	0.440	0.434	0.95	0.87–1.05	0.320
5q34	rs6556756	163821858	—	T/G	0.089	0.112	0.116	0.096	0.97	0.83–1.13	0.722
6p22.3	rs7754840	20769229	M	C/G	0.324	0.332	0.306	0.317	1.05	0.95–1.17	0.391
6q25.1	rs6922269	151294678	M	A/G	0.255	0.307	0.247	0.250	1.12	1.00–1.25	0.033
8q24.11	rs13266634	118253964	M	C/T	0.686	0.693	0.696	0.701	1.03	0.93–1.15	0.597
8q24.21	rs6983267	128482487	C	G/T	0.528	0.526	0.525	0.516	0.98	0.87–1.08	0.606
8q24.21	rs7014346	128493974	C	A/G	0.386	0.376	0.369	0.352	0.95	0.86–1.05	0.227
8q24.21	rs1447295	128554220	C	A/C	0.142	0.110	0.116	0.122	0.91	0.79–1.06	0.235
9p21.3	rs564398	22019547	M	T/C	0.572	0.559	0.596	0.563	0.90	0.82–1.00	0.037
9p21.3	rs10757278	22114477	M	G/A	0.457	0.461	0.476	0.428	0.90	0.81–0.99	0.026
9p21.3	rs1333049	22115503	M	C/G	0.542	0.543	0.527	0.570	1.11	1.00–1.22	0.030
9p21.3	rs10811661	22124094	M	T/C	0.823	0.808	0.796	0.826	1.08	0.96–1.23	0.216
10q23.33	rs1111875	94452862	M	C/T	0.593	0.594	0.598	0.605	1.02	0.93–1.13	0.680
10q25.2	rs7903146	114748339	M	T/C	0.278	0.275	0.264	0.273	1.02	0.92–1.14	0.720
10q26.13	rs2420946	123341314	C	T/C	0.392	0.359	0.396	0.391	0.93	0.84–1.103	0.167
11p15.1	rs5219	17366148	M	T/C	0.378	0.344	0.370	0.359	0.91	0.83–1.01	0.080
12q21.1	rs1495377	69863368	M	G/C	0.505	0.528	0.494	0.519	1.10	1.00–1.22	0.046
15q25.1	rs8034191	76593078	C	T/C	0.682	0.682	0.662	0.683	1.06	0.95–1.18	0.270
16q12.1	rs8051542	51091668	C	T/C	0.730	0.730	0.756	0.715	0.88	0.79–0.98	0.023
16q12.1	rs12443621	51105538	C	G/A	0.425	0.439	0.421	0.446	1.09	0.99–1.20	0.083
16q12.2	rs8050136	52373776	M	A/C	0.383	0.371	0.391	0.388	0.97	0.88–1.08	0.599
17q12	rs757210	33170628	M, C	T/C	0.384	0.381	0.387	0.384	0.99	0.90–1.09	0.772
17q12	rs4430796	33172153	M, C [†]	A/G	0.486	0.501	0.484	0.494	1.05	0.96–1.16	0.320
18q21.1	rs4939827	44707461	C	T/C	0.494	0.528	0.491	0.494	1.06	0.97–1.17	0.202
Xp11.22	rs5945572	51246423	C	A/G	0.354	0.375	0.341	0.366	0.99	0.91–1.08	0.852

*M indicates that the risk allele contributes to metabolic disease (coronary artery disease or type 2 diabetes); C indicates that the risk allele contributes to cancer.

[†]Major allele (A) has been associated with risk for cancer, whereas the minor allele (G) has been associated with type 2 diabetes.

[‡]Major allele is indicated in bold.

^sEstimation of OR and 95% CI was performed with robust SEs to take into account family dependency in the LLS.

^a*P* values were calculated using robust SEs to take into account family dependency in the LLS.

Table S6. Sum of risk alleles per age group

Age group	Age range	N	Mean sum of risk alleles (SE)					
			All		Metabolic		Cancer	
1	<26 y	116	26.56	(0.30)	18.68	(0.27)	9.70	(0.23)
2	26–35 y	576	26.80	(0.14)	18.62	(0.12)	10.01	(0.09)
3	36–45 y	247	26.72	(0.21)	18.71	(0.18)	9.79	(0.13)
4	46–55 y	344	26.92	(0.17)	18.58	(0.14)	10.08	(0.12)
5	56–65 y	457	27.11	(0.16)	18.60	(0.14)	10.09	(0.10)
6	66–75 y	135	26.75	(0.30)	18.25	(0.24)	10.17	(0.19)
7	76–85 y	547	26.83	(0.14)	18.63	(0.12)	10.01	(0.09)
8	86–95 y	1,029	26.78	(0.10)	18.61	(0.09)	9.92	(0.07)
9	96–98 y	101	26.82	(0.31)	18.25	(0.28)	10.11	(0.20)
10	≥99 y	38	26.84	(0.47)	19.61	(0.45)	9.29	(0.32)

Other Supporting Information Files

Dataset S1 (XLS)